

Brief Mindfulness Meditation for Depression and Anxiety Symptoms in Patients Undergoing Dialysis: *A Randomized Control Trial*



Zoë Thomas PGY-5
McGill Psychiatry
EAPM - 2017

Study Summary



- RCT set in urban dialysis unit
- 41 dialysis patients with depression/anxiety symptoms
- Randomly assigned to intervention (n=21) or TAU (n=20)
- 8 weeks, 3 times a week, 10-15 minutes chairside meditation
- Outcomes
 - 1^o: Feasibility – enrolment rates, completion rates, tolerability
 - 2^o: Depression & anxiety scores – PHQ-9 & GAD-7

Premise



- Depression & anxiety symptoms affect 50% of dialysis patients
 - Lower quality of life, increased dialysis non-adherence, suicidality, medical comorbidity and mortality
- Management generally inadequate
 - Low detection rates, limited access to psychotherapy, polypharmacy, minimal literature on effective treatments
- Alternative, effective, scalable strategies needed

Mindfulness Meditation



**Moment-to-moment
Non-judgmental
Awareness**

- Effective in several psychiatric conditions
 - Depression, anxiety, SUD
- Effective in several chronic physical illnesses
 - Psoriasis, breast/gyne cancer, IBD, DM, CAD, transplant
- Not tested in dialysis population

Methods



- Recruitment at Jewish General Hospital dialysis unit – 2016
- Inclusion criteria:
 - English or French
 - Scores of ≥ 6 on PHQ-9 and/or GAD- 7 scales
- Exclusion criteria:
 - Cognitive impairment (abnormal Mini-Cog)
 - Current psychosis
 - Acute suicidal ideation with intent
- Randomized 1:1


Intervention



- Chairside meditations 3 times a week, 10-15 minutes, for 8 weeks
 - *Body scan*
 - *Guided meditation*
 - *Silent meditation*
 - *Gentle arm movements*
- Brief check-ins before & after
- Home practice encouraged

Both groups received psychoeducational literature on depression & anxiety & local resources

Results



Results



Participant Data	Total Sample (n=41) Mean±SD	Intervention (n=21) Mean±SD	TAU (n=20) Mean±SD
<i>Demographic Information</i>			
Female	33 %	33 %	35 %
Age	65 ± 13	66 ± 13	64 ± 1 4
Married/common law	49 %	43 %	55 %
Non-White	51 %	48 %	55 %
Living with family	50 %	50 %	50 %
University	37 %	38 %	35 %
<i>Medical History</i>			
Number of medical problems	10 ± 4	11±4	10±4
Hypertension	85 %	91 %	80 %
Diabetes	63 %	52 %	75 %
Coronary artery disease	44 %	52 %	35 %
CHF / arrhythmia / PVD - each	32 %	43 %	20 %
Respiratory illness	29 %	33 %	25 %
Number of current medications	13 ± 5	14±4	12±5
Psychotropic medications	46 %	48 %	45 %
<i>Habits</i>			
Smoking	12 %	14 %	10 %
Alcohol /drugs	9 %	10 %	10 %

Results - Feasibility



- 71% retention rate
 - Mean number of sessions = 20/24 (82% of sessions)
 - Home practice average = 3 days a week x 14 minutes
- Meditation tolerability **8/10**
 - Guided meditation > gentle arm movements
- No adverse events
 - 10 participants admitted to hospital

Results - Effectiveness



Measure	Intervention Group			TAU Group			Stats	
	Pre n=21	Post n=17	Change n=17	Pre n=20	Post n=15	Change n=15	p	d
PHQ-9	12.7 ± 4.2	9.6 ± 4.9	-3.0 ± 3.9	11.9 ± 5.8	7.8 ± 5.1	-2.0 ± 4.7	p=0.45	0.13
GAD-7	6.2 ± 5.1	6.1 ± 4.7	-0.9 ± 4.6	6.1 ± 5.8	3.5 ± 2.1	-0.8 ± 4.8	p=0.91	0.35

Subjective Reports

Courage, Hope, Dignity,
Self-confidence, Strength

Discussion



- Underpowered?
- Inclusion of milder psychiatric symptomatology?
- Inclusion of severely medically ill patients?
- Lack of cognitive component of intervention?
- Lack of qualitative questionnaires?

Conclusion



Need for depression & anxiety treatments in dialysis patients

Barriers to usual treatments

- Additional appointments
- Polypharmacy
- Limited effectiveness

Meditation seems promising

- Scalable
- Feasible
- Enjoyable

Recommend larger studies

- N=60
- PHQ-9 > 10
- Qualitative scales
- Cognitive component

References



- Zalai D, Szeifert L, Novak M. Psychological distress and depression in patients with chronic kidney disease. *Semin Dial.* 2012;25(4):428-38.
- Cukor D, Coplan J, Brown C, Friedman S, Cromwell-Smith A, Peterson RA, Kimmel PL. Depression and anxiety in urban hemodialysis patients. *Clin J Am Soc Nephrol.* 2007 May;2(3):484-90. Epub 2007 Apr 4.
- Olagunju AT, Campbell EA, Adeyemi JD. Interplay of anxiety and depression with quality of life in endstage renal disease. *Psychosomatics.* 2015;56(1):67-77.
- Kovacs AZ, Molnar MZ, Szeifert L, Ambrus C, Molnar-Varga M, Szentkiralyi A, et al. Sleep disorders, depressive symptoms and health-related quality of life--a cross-sectional comparison between kidney transplant recipients and waitlisted patients on maintenance dialysis. *Nephrol Dial Transplant.* 2011;26(3):1058-65.
- Novak, M, et al., Depressive symptoms and mortality in patients after kidney transplantation: a prospective prevalent cohort study. *Psychosom Med.* 2010. 72(6): p. 527-34.
- García-Llana H, Remor E, Del Peso G, Selgas R. The role of depression, anxiety, stress and adherence to treatment in dialysis patients' health-related quality of life: a systematic review of the literature. *Nefrologia.* 2014 Sep 26;34(5):637-57
- Pompili M, Venturini P, Montebovi F, Forte A, Palermo M, Lamis DA, Serafini G, Amore M, Girardi P. Suicide risk in dialysis: review of current literature. *Int J Psychiatry Med.* 2013;46(1):85-108.
- Farrokhi F. Depression among dialysis patients: barriers to good care. *Iran J Kidney Dis.* 2012;6(6):403-6.
- Palmer SC1, Natale P, Ruospo M, Saglimbene VM, Rabindranath KS, Craig JC, Strippoli GF. Antidepressants for treating depression in adults with end-stage kidney disease treated with dialysis. *Cochrane Database Syst Rev.* 2016;(5):CD004541. doi: 10.1002/14651858.CD004541.pub3.
- Kabat-Zinn J, *Clinic UoMMCWSR.* Full catastrophe living: using the wisdom of your body and mind to face stress, pain, and illness: Pub. by Dell Publishing, a division of Bantam Doubleday Dell Pub. Group; 1990.

References



- Jacquart J, Miller KM, Radossi A, Haime V, Macklin E, Gilburd D, et al. The effectiveness of a community-based, mind-body group for symptoms of depression and anxiety. *Adv Mind Body Med.* 2014;28(3):6-13.
- Miller KM, Chad-Friedman E, Haime V, Mehta DH, Lepoutre V, Gilburd D, et al. The effectiveness of a brief mind-body intervention for treating depression in community health center patients. *Glob Adv Health Med.* 2015;4(2):30-5.
- Zgierska A, Rabago D, Chawla N, Kushner K, Koehler R, Marlatt A. Mindfulness meditation for substance use disorders: a systematic review. *Subst Abus.* 2009;30(4):266-94.
- Linda E. Carlson. Mindfulness-Based Interventions for Physical Conditions: A Narrative Review Evaluating Levels of Evidence. *International Scholarly Research Network ISRN Psychiatry.* 2012: 2012: 1-21.
- Kabat-Zinn J, Wheeler E, Light T, Skillings A, Scharf MJ, Cropley TG, Hosmer D, Bernhard J. Influence of a Mindfulness Meditation-Based Stress Reduction Intervention on Rates of Skin Clearing in Patients With Moderate to Severe Psoriasis Undergoing Photo Therapy (UVB) and Photochemotherapy (PUVA). *Psychosomatic Medicine.* 1998;60(5): 625-632.
- Kuo B, Bhasin M, Jacquart J, Scult MA, Slipp L, Riklin EI, et al. Genomic and clinical effects associated with a relaxation response mind-body intervention in patients with irritable bowel syndrome and inflammatory bowel disease. *PLoS One.* 2015;10(4):e0123861
- Gerbarg PL, Jacob VE, Stevens L, Bosworth BP, Chabouni F, DeFilippis EM, et al. The Effect of Breathing, Movement, and Meditation on Psychological and Physical Symptoms and Inflammatory Biomarkers in Inflammatory Bowel Disease: A Randomized Controlled Trial. *Inflamm Bowel Dis.* 2015;21(12): 2886-2896.
- Bower JE, Crosswell AD, Stanton AL, Crespi CM, Winston D, Arevalo J, Ma J, Cole SW, Ganz PA. Mindfulness Meditation for Younger Breast Cancer Survivors: A Randomized Controlled Trial. *Cancer* 2015;121(8):1231-1241

References



- Keyworth C, Knopp J, Roughley K, Dickens C, Bold S, Coventry P. A Mixed-Methods Pilot Study of the Acceptability and Effectiveness of a Brief Meditation and Mindfulness Intervention for People with Diabetes and Coronary Heart Disease. *Journal of Behavioral Medicine*. 2014;40(2): 53-64.
- Watnick S, Wang PL, Dematura T, Ganzini L. Validation of 2 Depression Screening Tools in Dialysis Patients. *Am J Kidney Dis*. 2005; 46(5): 919-924.
- Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*. 2001;16(9):606-13.
- Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. 2006;166(10):1092-7.
- Segal ZV, Williams JMG, Teasdale JD. *Mindfulness-Based Cognitive Therapy for Depression, Second Edition*. 2013. Guildford Press: New York
- Barrett MS, Chua W-J, Crtis-Christoph P, Gibbons MB, Casiano D, Thompson D. Early withdrawal from mental health treatment: implications for psychotherapy practice. *Psychotherapy*. 2008;45(2):247-267.
- Cukor D, Ver Halen N, Asher DR, Coplan JD, Weedon J, Wyka KE, Saggi SJ, Kimmel PL. Psychosocial intervention improves depression, quality of life, and fluid adherence in hemodialysis. *J Am Soc Nephrol*. 2014;25(1):196-206.
- Duarte PS, Miyazaki MC, Blay SL, Sesso R. Cognitive-behavioral group therapy is an effective treatment for major depression in hemodialysis patients. *Kidney International*. 2009;76(4): 414-421.
- Erdley SD, Gellis ZD, Bogner HA, Kass DS, Green JA, Perkins RM. Problem-solving therapy to improve depression scores among older hemodialysis patients: a pilot randomized trial. *Clinical Nephrology*. 2014;82(1):26-33.
- Vamos M. Psychotherapy in the medically ill: A Commentary. *Aust N Z J Psychiatry*. 2006 April; 40(4): 295-309.
- Haugan G, Utvaer BK, Moksnes UK. The Herth Hope Index – A psychometric study among cognitively intact nursing home patients. *J Nurs Meas*. 2013;21(3):378-400.



Thank you



zoe.thomas@mail.mcgill.ca